
Acute Meningitis: Diagnosis, Interpretation, & Controversy

Heather Prendergast, MD, MPH, FACEP



2009 MEMC V Meeting



Valencia, Spain
16 September 2009

Heather Prendergast, MD, MPH, FACEP



**Heather M. Prendergast, MD,
MPH, FACEP**

Associate Professor

Department of Emergency Medicine
University of Illinois College of Medicine
Chicago, IL

Heather Prendergast, MD, MPH, FACEP



Disclosures

- FERNE Executive Board
- FERNE grant support by industry

Heather Prendergast, MD, MPH, FACEP



Objectives

- Present a relevant patient case
- Discuss contraindications for lumbar puncture (LP)
- Review the procedure of LP
- Discuss the role of antibiotics , anti-virals, and steroids.

Heather Prendergast, MD, MPH, FACEP




A Clinical Case

Heather Prendergast, MD, MPH, FACEP



ED Presentation


- 77 yo previously healthy female
- 3 day history of fever, confusion, and lethargy
- Glasgow Coma Scale 13 (E4,V4,M5)
- Key Aspects of Physical Exam:
 - Unable to cooperate with neurological exam, +neck stiffness upon neck flexion

Heather Prendergast, MD, MPH, FACEP 

Indications for pre-LP head CT scan


Table 2. Recommended criteria for adult patients with suspected bacterial meningitis who should undergo CT prior to lumbar puncture (B-II).

Criterion	Comment
Immunocompromised state	HIV infection or AIDS, receiving immunosuppressive therapy, or after transplantation
History of CNS disease	Mass lesion, stroke, or focal infection
New onset seizure	Within 1 week of presentation; some authorities would not perform a lumbar puncture on patients with prolonged seizures or would delay lumbar puncture for 30 min in patients with short, convulsive seizures
Papilledema	Presence of venous pulsations suggests absence of increased intracranial pressure
Abnormal level of consciousness	---
Focal neurologic deficit	Including dilated nonreactive pupil, abnormalities of ocular motility, abnormal visual fields, gaze palsy, arm or leg drift


Heather Prendergast, MD, MPH, FACEP 

Contraindications


- Skin infection near site of LP
- Suspicion of increased intracranial pressure due to cerebral mass
- Uncorrected coagulopathy
- Acute spinal cord trauma

Heather M. Prendergast, MD, MPH 


Positioning



INCORRECT




CORRECT

Heather Prendergast, MD, MPH FACEP 

Predicting difficult and traumatic lumbar punctures.


The American Journal of Emergency Medicine
 2007, Volume 25, Issue 6, Pages 608-611
 K. Shah, D. McGillicuddy, J. Spear, J. Edlow

- Difficult LP
 - Requires 3 or more needle sticks or attempt by another clinician
- Spine visibility
 - Ability to see the contour of the spinous processes
- Spine palpability
 - Ability to palpate distinct spinous processes

Heather Prendergast, MD, MPH FACEP 

Comparison of Clinical Groups

	Difficult	Traumatic
• Spine visible	20.0	8.6
• Spine not visible	42.0	21.8
• Spine palpable	26.7	12.4
• Spine not palpable	44.2	23.3
• BMI > 30	42.1	18.4
• BMI < 30	28.2	14.6
• Age >65	42.9	14.3
• Age <65	29.9	15.8

Heather Prendergast, MD, MPH FACEP 

Ultrasound Assisted Lumbar Puncture

- 46 patients
 - 22 Palpation Landmarks (PL)
 - 24 Ultrasound Landmarks (UL)
- Failure Rates
 - 6/22 PLs
 - 1/24 ULs
- Obese Patients
 - 4/7 failed PLs
 - 0/5 failed ULs

Heather Prendergast, MD, MPH, FACEP



Understanding Opening Pressures

- Normal: 60-200 mm H₂O (obese patients up to 250mm H₂O)
- Elevated: Suggest increased intracranial pressures (>250 mm H₂O)
 - Mass lesion (neoplasm, hemorrhage, infection)
 - Overproduction of CSF
 - Defective Outflow Mechanics

Heather M. Prendergast, MD, MPH



CSF Composition

- Color
 - Clear and colorless
 - Turbid
 - 200 WBCs or 400 RBCs
 - Grossly Bloody
 - 6000 RBCs

Heather Prendergast, MD, MPH, FACEP



CSF Composition

- Cells
 - Acellular (up to 5 WBCs and 5 RBCs)
 - More than 3 polymorphonuclear leukocytes (PMNs) abnormal

Heather Prendergast, MD, MPH, FACEP



CSF Pleocytosis

- CSF pleocytosis
 - 10 white blood cells/ μ L, corrected for CSF red blood cells using a ratio of 1 WBC per 500 RBCs

Heather M. Prendergast, MD, MPH



Calculating Predicted CSF WBC count

Predicted CSF WBC count/ μ L =

CSF RBC count X (peripheral blood WBC count \div peripheral RBC count)

Heather M. Prendergast, MD, MPH



Validation of Prediction Calculation in Adults

- 720 patients
 - CSF WBC count >10X predicted value
 - Positive Predictive Value 48% for Bacterial Meningitis
 - CSF WBC count < 10X predicted value
 - Negative Predictive Value 99% for meningitis

Heather M. Prendergast, MD, MPH



Validation of Prediction Calculation in Children

- 92 children
 - CSF WBC count >10X predicted value
 - 28/30 children (93%) bacterial meningitis
 - 57 children
 - CSF WBC count < 10X predicted
 - 100% for predicting the absence of meningitis

Heather M. Prendergast, MD, MPH



CSF Composition

- Protein
 - Largely excluded from CSF by blood-CSF barrier
 - Normal range (adults) 23-38 mg/dL
 - False elevation
 - Diabetes, Presence of RBCS
 - True elevation
 - Infectious and Noninfectious Conditions

Heather M. Prendergast, MD, MPH



CSF Composition

- Glucose
 - CSF-to-serum glucose ratio
 - Normal 0.6
 - Low CSF glucose concentrations
 - Bacterial meningitis
 - Mycobacterial and Fungal CNS infections
 - M. pneumoniae and Noninfectious processes
 - Less than 18 mg/dL strongly predictive of bacterial meningitis

Heather Prendergast, MD, MPH, FACEP



CSF Composition

- Glucose
 - CSF-to-serum glucose ratio
 - Limited utility in Neonates, and severe hyperglycemia
 - Normal CSF glucose concentrations
 - Viral CNS infections
 - Exceptions:
 - mumps, enteroviruses, lymphocytic choriomeningitis(LCM), herpes simplex

Heather Prendergast, MD, MPH, FACEP



CSF Composition

- Lactate
 - Elevated in bacterial meningitis
 - One study higher sensitivity and specificity than blood glucose ratio

Heather Prendergast, MD, MPH, FACEP



CSF in CNS Infection

- Bacterial Meningitis
 - CSF WBC > 1000/microL (with PMNs)
 - CSF Protein >250 mg/dL
 - CSF Glucose < 45 mg/dL (2.5 mmol/L)
 - CSF-blood glucose ratio 0.4 or less (LR 18)
 - CSF Lactate >31.53 mg/dL(3.5 mmol/L)

Heather Prendergast, MD, MPH, FACEP



CSF in CNS Infection

- Viral Meningitis
 - CSF WBC < 250 /microL (with lymphocytes)
 - CSF Protein <150 mg/dL
 - CSF Glucose more than 50% of serum concentration

Heather Prendergast, MD, MPH, FACEP



Summary of Typical CSF Findings

	Normal	Bacterial	Viral	Fungal
Opening Pressure	<170 mm	>300	200	300
Cells	0-5	>1000	<1000	< 500
Polymorphs	0	Predominate	Early	+/- increased
Lymphocytes	5	Late	Predominate	Increased
Glucose	60-80	<40	>40	<40
CSF plasma: Glucose ratio	66%	<40%	Normal	< 30%
Protein	5-40	Increased >200	+/- Increased <200	Increased >200
Culture	Negative	Positive	Negative	+positive

Heather Prendergast, MD, MPH, FACEP



Organisms

- < 3 months • *E.Coli , Listeria, Streptococci*
- 3 months- 18 yrs • *N. meningitidis, H. influenzae, S. pneumoniae*
- 18yrs – 50 yrs. • *N. meningitidis, S. pneumoniae*
- > 50 yrs. • *S. pneumoniae, Listeria, gram-negative bacilli*

Heather Prendergast, MD, MPH, FACEP



Antibiotic Choices (Recommendations)

- Good CNS penetration
 - Ceftriaxone
 - Cefotaxime
 - Vancomycin (controversial in children)
- Listeria (young, old, immunosuppressed)
 - Add high-dose ampicillin
- PCN or Cephalosporin Allergy
 - Meropenem or Chloramphenicol + Vancomycin

Heather M. Prendergast, MD, MPH



PEDIATRICS

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Early Vancomycin Therapy and Adverse Outcomes in Children With Pneumococcal Meningitis

PEDIATRICS Vol. 117 No. 5 May 2006, pp. 1688-1694

- Associated with a substantially increased risk of hearing loss.
- Recommend delaying the first dose of vancomycin therapy until 2 hours after the first dose of parenteral cephalosporin

Heather M. Prendergast, MD, MPH



Antivirals & Meningitis

- Viruses
 - Arbovirus
 - Herpes Simplex *
 - Cytomegalovirus
 - Adenovirus
 - HIV
- * Protracted course (Acyclovir 10mg/kg IV Q8 hours)

Heather M. Prendergast, MD, MPH



Controversy Steroid Use In Bacterial Meningitis

Heather Prendergast, MD, MPH, FACEP



Steroids & Meningitis

European Journal of Neurology 2009, 16: 662-673

doi:10.1111/j.1468-1331.2009.02815.x

REVIEW ARTICLE

Adjunctive dexamethasone therapy for bacterial meningitis in adults: a meta-analysis of randomized controlled trials

K. Z. Vardakas^a, D. K. Matthaiou^a and M. E. Falagas^{a,b}

^aAlfa Institute of Biomedical Sciences (AIBS), Athens, Greece; and ^bDepartment of Medicine, Tufts University School of Medicine, Boston, MA, USA

Heather M. Prendergast, MD, MPH




Meta-Analysis & Malawian RCT

Analyses with the inclusion of the Malawian RCT

Mortality, prior antibiotic use (0.79, 0.45-1.37)
 Mortality, no prior antibiotic use (0.68, 0.36-1.28)
 Mortality, symptoms < 48 h before hospitalization (0.61, 0.36-1.00)
 Mortality, symptoms > 48 h before hospitalization (0.80, 0.47-1.36)
 Mortality, definite meningitis (0.55, 0.31-0.96)
 Mortality, double blind, (0.87, 0.66-1.14)
 Unfavorable outcome (0.78, 0.54-1.13)

Analyses without the inclusion of the Malawian RCT

Mortality, prior antibiotic use (0.68, 0.43-1.07)
 Mortality, no prior antibiotic use (0.44, 0.24-0.82)
 Mortality, symptoms < 48 h before hospitalization (0.45, 0.23-0.87)
 Mortality, symptoms > 48 h before hospitalization (0.66, 0.43-1.03)
 Mortality, definite meningitis (0.41, 0.26-0.66)
 Mortality, double blind (0.60, 0.39-0.93)
 Unfavorable outcome (0.70, 0.44-1.12)



Heather M. Prendergast, MD, MPH



Meta-Analysis Results

- Treatment with Dexamethasone Benefits:
 - Lower mortality than Placebo (non-significant OR= 0.58)
 - Lower mortality with definite meningitis (OR=0.55)
 - Shorter duration of symptoms (OR=0.61)
 - *Streptococcus Pneumoniae* meningitis (OR=0.26)

Heather M. Prendergast, MD, MPH



Meta-Analysis Conclusions

- Treatment with Dexamethasone
 - Did not decrease mortality in all patients
 - Beneficial in the following groups
 - Early presentation (< 48 hours)
 - Definite meningitis
 - *Streptococcus Pneumoniae* meningitis
 - Countries with medium or high HDI

Heather M. Prendergast, MD, MPH




Disponible en ligne sur ScienceDirect ELSEVIER MASSON
www.sciencedirect.com
EM|consulte www.em-consulte.com
Médicine et maladies infectieuses 39 (2009) 531-538

Texts of experts
Corticosteroids for acute adult bacterial meningitis[☆]
Corticothérapie pour les méningites bactériennes de l'adulte
D. van de Beek
Department of Neurology, Centre of Infection and Immunity of Amsterdam (CIIMA), Academic Medical Centre, University of Amsterdam, P.O. Box 22700, 1100 Amsterdam, The Netherlands
Received 15 January 2009; accepted 20 February 2009
Available online 21 April 2009

- Dexamethasone given before or with 1st dose of antibiotics
- No serious adverse effects
- Consistent beneficial effect on mortality and borderline statistical beneficial effect on neurologic sequels

Heather M. Prendergast, MD, MPH



Corticosteroids for Acute Bacterial Meningitis

Ann Emerg Med. 2009 Jul;54(1):136-7

To the Editor:

We are writing in response to the systematic review abstract entitled “Corticosteroids for Acute Bacterial Meningitis”

However, we disagree with some of the conclusions:

Both sets of authors, as a result of the review, advocate corticosteroids for all adults and children with suspected bacterial meningitis. While undoubtedly correct in the era before the H. influenzae vaccine, we feel this recommendation may not be the current best course of action for pediatric patients.

Heather M. Prendergast, MD, MPH



Case Resolution

- CT scan: No mass lesion
 - CSF Results
 - WBC 5000 / μ L
 - RBC 5 /microL
 - CSF blood glucose ratio 0.2
 - Gram stain: gram positive rods
- Diagnosis: Meningitis due to Listeria

Heather Prendergast, MD, MPH, FACEP



Recommendations

- Calculate CSF-blood glucose ratio.
 - 0.4 or less (LR 18) bacterial meningitis
- Determine the predicted CSF WBC count
 - Negative Predictive Value 99% for bacterial meningitis

Heather Prendergast, MD, MPH, FACEP



Recommendations

- Do not delay administering antibiotics as delays have been associated with worsening clinical outcomes
- Empiric antibiotic choice is based upon broad spectrum coverage of common pathogens
- Use of dexamethasone has shown benefit in selected cases of adult bacterial meningitis less so in children

Heather Prendergast, MD, MPH FACEP



Conclusions

- Primary indications for LP is to assess for meningitis or subarachnoid hemorrhage
- Most patients do not require CT scan to rule out mass lesions prior to LP
- Clinical examination can guide decision for neuroimaging
- Elevated opening pressures indicate increase intracranial pressures

Heather Prendergast, MD, MPH FACEP



Questions?

hprender@uic.edu

Heather Prendergast, MD, MPH, FACEP

